Centrum voor Wiskunde en Informatica

REPORT RAPPORT

Homogeneous behaviors

M.S. Ravi, J. Rosenthal and J.M. Schumacher

Department of Operations Reasearch, Statistics, and System Theory

BS-R9607 1996

Report BS-R9607
ISSN 0924-0659

CWI
P.O. Box 94079
1090 GB Amsterdam
The Netherlands

Homogeneous Behaviors

M.S. Ravi
Department of Mathematics, East Carolina University
Greenville, North Carolina 27858, USA
maravi@ecuvax.cis.ecu.edu

J. Rosenthal*
Department of Mathematics, University of Notre Dame
Notre Dame, Indiana 46556, USA
Joachim.Rosenthal@nd.edu

J.M. Schumacher
CWI
P.O. Box 94079, 1090 GB Amsterdam, The Netherlands, and
Tilburg University, CentER and Department of Economics,
P.O. Box 90153, 5000 LE Tilburg, The Netherlands
Hans.Schumacher@cwi.nl

Abstract

Recently a smooth compactification of the space of linear systems with n states, m inputs and p outputs has been discovered. In this paper we obtain a concrete interpretation of this compactification as a space of discrete-time behaviors. We use both homogeneous polynomial representations and generalized first-order representations, and provide a realization theory to link these to each other.

AMS Subject Classification (1991): 93B20, 93B27.
Keywords & Phrases: generalized linear systems, smooth compactification, discrete-time behaviors.
Note: This report has been submitted for publication elsewhere.

1 Introduction

An analogy that has been instrumental in the development of the 'behavioral' theory of linear systems [14, 15, 16] is that between linear behaviors and linear subspaces of $\mathbb{R}^n$. In fact a subspace of $\mathbb{R}^n$ can be looked at as a static behavior, and the theory of linear time-invariant behaviors may be viewed as the most direct dynamic generalization of it. The analogy makes it natural for instance to look for 'kernel' and 'image' representations (corresponding to 'AR' and 'MA' representations in the terminology of [16]). It is also suggestive in defining a notion of convergence. Starting from the representation of a subspace of given dimension m as the column space of a matrix of size $n \times m$, which is determined up to right multiplication by nonsingular matrices, a natural notion of convergence for subspaces is obtained by the construction of the quotient topology. One may also use the kernel representation in the same way; fortunately, this leads to the same notion of convergence. In the same manner, one can define a topology for the the set of linear time-invariant systems of a given state space dimension n and a given number of inputs m. Starting for instance from a minimal first-order representation, which is unique up to similarity transformations, a notion of convergence is again provided by the

*This author was supported in part by NSF grant DMS-94-00965. The research for this paper was carried out in part while he was a visitor at CWI.

quotient topology. The set of m-dimensional subspaces of $\mathbb{R}^n$, with the indicated topology, is usually denoted by Grass(m,n); we shall use the symbol Sys(m,p,n) to indicate the set of time-invariant linear systems with m inputs, p outputs, and an n-dimensional state space.

In spite of the analogy between 'subspaces' and 'systems', there is an important topological difference between Grass(m,n) and Sys(m,p,n). Whereas the set of subspaces is compact, so that every sequence must have a limit point, the set of systems is not compact. To give an example in the very simple case $n=1$, $m=0$, $p=1$, the discrete-time behavior spanned by the sequence $(1,a^{-1},a^{-2},\cdots)$ does not converge to a limit if a tends to zero. (Compare this to the sequence of one-dimensional subspaces spanned by the vector $(1,a^{-1},\cdots,a^{-n})$; as a tends to 0, this sequence does converge to a limit, namely the subspace spanned by $(0,0,\cdots,1)$.) A limit behavior can only be obtained if one enlarges the space of possible behaviors. In our example, a suggestion of how to do this is arrived at as follows. Identify a sequence $w=(w_1,w_2,\ldots)$ with the 'one-sided' power series $w(s):=\sum_{i=1}^{\infty} w_i s^{-i}$. In this way w_a is identified with the power series

$$w_a(s) := \sum_{i=1}^{\infty} a^{-i+1} s^{-i} = \frac{a}{as-1}. \tag{1.1}$$

Given that the meromorphic function $(as-1)^{-1}$ converges to the constant function -1 uniformly on compact subsets of $\mathbb{C}$ as a tends to zero, a natural candidate for a limit point of the one-dimensional subspaces generated by $w_a(s)$ would be the subspace generated by $w_0(s)=1$. Note that this subspace is not generated by a power series of the form $w(s)=\sum_{i=1}^{\infty} w_i s^{-i}$. The example suggests that one somehow should add some 'generalized systems' which could serve as additional limit points.

It was recently discovered that it is possible to compactify the set Sys(m,p,n) in a "smooth manner" using either kernel representations by homogeneous polynomial matrices [10] or first-order representations under weakened minimality conditions [3]. A continuous-time interpretation of generalized systems in terms of impulsive-smooth behaviors was given in [4], and a discrete-time interpretation on a rather abstract level has been provided in [9]. We still believe that is is of interest to add an interpretation in terms of concrete discrete-time behaviors, since the availability of an interpretation that is as simple and as concrete as possible may contribute to the understanding of the compactified space of systems.

The purpose of the present paper is therefore to obtain a behavioral interpretation in discrete time of the systems given by homogeneous higher-order equations as in [10] or by generalized first-order equations as in [4]. The resulting set of behaviors, which is slightly bigger than the set considered by J.C. Willems for instance in [14], will be referred to as the set of 'homogeneous behaviors'.

The organization of this paper is as follows. The framework that we shall use will be introduced in Section 2. Here we also introduce the various representations of homogeneous behaviors. The relations between these representations will be explored in some detail in Section 3. In this section we also describe a very simple realization algorithm for homogeneous autoregressive systems first announced in [12]. In Section 4 we obtain the main result of the paper, establishing a one-to-one connection between (i) homogeneous polynomial matrices modulo homogeneous unimodular row transformations, (ii) generalized first-order representations modulo similarity transformations, and (iii) homogeneous behaviors. Conclusions follow in Section 5.

2 DEFINITIONS AND PRELIMINARIES

Denote by $\mathcal{L}$ the set of all functions from the integers $\mathbb{Z}$ to the reals $\mathbb{R}$ with support bounded on the left; that is, $\mathcal{L}$ is the set of two-sided infinite sequences of real numbers $(\cdots,a_{-1},a_0,a_1,\cdots)$ for which there exists an N such that $a_i=0$ for all $i<-N$. Of course, $\mathbb{Z}$ can be thought of as a time axis and $\mathcal{L}$ as a signal space. We shall write $\mathcal{L}^q$ for the analogous space of sequences with values in $\mathbb{R}^q$; in the terminology of [14], this will be our *universum*.

We shall use two notations for the elements of $\mathcal{L}^q$. The first one is the sequence notation

$$w \;=\; (w_{-N},\cdots,w_{-1},w_0 \mid w_1,w_2,\cdots) \tag{2.1}$$

where it is understood that $w_i = 0$ for $i < -N$, and in which we use a vertical bar to indicate the position between the signal values at time points 0 and 1. The second notation is the representation as a formal Laurent series:

$$w(s) = \sum_{i=-N}^{\infty} w_i s^{-i}. \tag{2.2}$$

Such series may be thought of as consisting of a 'polynomial part' $w_+(s) := \sum_{i=-N}^{0} w_i s^{-i}$ and a 'strictly proper part' $w_-(s) := \sum_{i=1}^{\infty} w_i s^{-i}$, corresponding to the parts before and after the bar in the sequence notation. The Laurent series is written in powers of s^{-1} rather than in powers of s so that the polynomial part is a polynomial in s rather than in s^{-1}; of course this is just a matter of notation. We shall write $\mathcal{L}_+^q$ for the set of elements of $\mathcal{L}^q$ that appear as polynomials in the Laurent series notation, and the set of elements whose polynomial part is zero will be denoted by $\mathcal{L}_-^q$. We remark that the spaces $\mathcal{L}^q$, $\mathcal{L}_+^q$, $\mathcal{L}_-^q$ have been well studied in the systems literature and we refer e.g. to [6].

Elements of $\mathcal{L}$ can be multiplied in the way suggested by the Laurent series notation (2.2). With respect to this multiplication and the usual addition, the set $\mathcal{L}$ is a *field* (compare with [6]) and we shall make use of this fact in this paper. Note that the field of rational functions $\mathbb{R}(s)$ can be viewed as a subfield of $\mathcal{L}$ by identifying $f(s) \in \mathbb{R}(s)$ with its Laurent expansion around infinity.

On $\mathcal{L}^q$ we shall consider four shift operators, namely left and right shifts with and without cancellation. In sequence notation, the four operators are defined by

$$\begin{array}{rl} \sigma: & (w_{-N},\cdots,w_{-1},w_0 \mid w_1,w_2,\cdots) \;\mapsto\; (w_{-N},\cdots,w_0,w_1 \mid w_2,w_3,\cdots) \\ \sigma_0: & (w_{-N},\cdots,w_{-1},w_0 \mid w_1,w_2,\cdots) \;\mapsto\; (w_{-N},\cdots,w_0,0 \mid w_2,w_3,\cdots) \\ \tau: & (w_{-N},\cdots,w_{-1},w_0 \mid w_1,w_2,\cdots) \;\mapsto\; (w_{-N},\cdots,w_{-2},w_{-1} \mid w_0,w_1,w_2,\cdots) \\ \tau_0: & (w_{-N},\cdots,w_{-1},w_0 \mid w_1,w_2,\cdots) \;\mapsto\; (w_{-N},\cdots,w_{-2},w_{-1} \mid 0,w_1,w_2,\cdots). \end{array} \tag{2.3}$$

The same operators can be given in Laurent series notation by

$$\begin{array}{rcl} (\sigma w)(s) & = & s w(s) \\ (\sigma_0 w)(s) & = & s w(s) - s w_-(s)|_{s=\infty} \\ (\tau w)(s) & = & s^{-1} w(s) \\ (\tau_0 w)(s) & = & s^{-1}(w(s) - w_+(0)). \end{array} \tag{2.4}$$

The following relations are easily seen to hold:

$$\sigma\tau = \tau\sigma = \text{identity} \tag{2.5}$$

and for all $k \geq 0$

$$\sigma_0^k \tau^k = \sigma^k \tau_0^k = \sigma_0^k \tau_0^k. \tag{2.6}$$

Consider now a homogeneous polynomial row vector of degree ν, with q entries. Such a row vector may be written in the form

$$p(s,t) = \sum_{k=0}^{\nu} p_k s^k t^{\nu-k}$$

where the p_k are constant row vectors of length q. We want to associate a linear operator on $\mathcal{L}^q$ to this homogeneous vector, which in turn will determine a 'behavior' as the set of all elements in $\mathcal{L}^q$ that are mapped to zero by this operator. Two linear operators that may be associated to $p(s,t)$ are the following:

$$p(\sigma_0,\tau) \overset{\text{def}}{=} \sigma_0^{\nu} \sum_{k=0}^{\nu} p_k \tau^{\nu-k} \tag{2.7}$$

and

$$p(\sigma,\tau_0) \stackrel{\text{def}}{=} \tau_0^\nu \sum_{k=0}^{\nu} p_k \sigma^k. \tag{2.8}$$

Both operators have the desirable property that, in the scalar case $q = 1$, the dimension of the associated behavior on $\mathcal{L}$ is equal to the degree of the polynomial from which the operator is derived. More is true: the following proposition shows that the behaviors determined by the two operators are in fact equal.

Proposition 2.1 *For any homogeneous polynomial $p(s,t)$, one has*

$$\{w \in \mathcal{L}^q \mid p(\sigma_0,\tau)w = 0\} \;=\; \{w \in \mathcal{L}^q \mid p(\sigma,\tau_0)w = 0\}. \tag{2.9}$$

Proof The statement follows from the identities

$$\sigma_0^\nu \sum_{k=0}^{\nu} p_k \tau^{\nu-k} \;=\; \sigma_0^\nu \tau^\nu \sum_{k=0}^{\nu} p_k \sigma^k \;=\; \sigma^\nu \tau_0^\nu \sum_{k=0}^{\nu} p_k \sigma^k$$

and from the fact that σ^ν is invertible. □

Remark 2.2 The proposition shows that from the point of view of behaviors it is immaterial whether one takes $p(\sigma_0,\tau)$ or $p(\sigma,\tau_0)$ as the operator associated to a homogeneous vector polynomial $p(s,t)$. One might say, though, that the choice $p(\sigma_0,\tau)$ is closer to tradition in the sense that it acts on one-sided sequences (interpreted as sequences in $\mathcal{L}^q$) in the same way as the standard operator associated to the dehomogenization $p(s) = p(s,1)$ of $p(s,t)$. In this context the effect of a factor t in $p(s,t)$ may be described as 'cancel one more element to the left'.

Remark 2.3 The association $p(s,t) \mapsto p(\sigma_0,\tau)$ is linear, but does not respect the multiplicative structure of homogeneous polynomials. For instance the operator associated to the homogeneous polynomial t^2 is not the square of the operator associated to t.

Remark 2.4 One might consider other sequence spaces than $\mathcal{L}^q$, which can be looked at as the space of bi-infinite sequences with support bounded to the left. In particular one might take the space of *all* bi-infinite sequences. It seems to be hard, though, to associate to homogeneous polynomials a linear operator on this space in such a way that the dimension of the kernel of this operator is equal to the degree of the homogeneous polynomial that one started with. If one associates to $p(s,t)$ the operator $p(\sigma_0,\tau)$ as defined above, then for instance the operator associated to $p(s,t) = s-t$, which is $\sigma_0(1-\tau)$, has a two-dimensional solution space associated to it (spanned by the sequences $(\cdots,1,1 \mid 0,0,\cdots)$ and $(\cdots,0,0 \mid 1,1,\cdots)$). On the other hand, if one would consider only sequences with finite support, then the same operator would have a zero-dimensional solution space. Finally, the use of sequences with support bounded to the right would give rise to a theory that is essentially the same as the one developed in this paper.

To a homogeneous polynomial matrix $P(s,t)$ we associate an operator $P(\sigma_0,\tau)$ by replacing each row $P_i(s,t)$ by the corresponding operator $P_i(\sigma_0,\tau)$. We now define:

Definition 2.5 The *homogeneous behavior* $\mathcal{B}(P)$ associated with a homogeneous polynomial matrix $P(s,t)$ is the set

$$\mathcal{B}(P) = \{w \in \mathcal{L}^q \mid P(\sigma_0,\tau)w = 0\}. \tag{2.10}$$

EXAMPLE 2.6 In continuation of the example discussed in the Introduction, consider the set of scalar homogeneous polynomials of degree 1, i. e. the set of polynomials $p_{a,b}(s,t) = as+bt$ with $(a,b) \neq (0,0)$. It is easily verified that for $a \neq 0$ the homogeneous behavior $\mathcal{B}(p_{a,b})$ is the one-dimensional space generated by the sequence $(0 \mid 1, -\frac{b}{a}, \frac{b^2}{a^2}, -\frac{b^3}{a^3}, \cdots)$, whereas for $a = 0$ the homogeneous behavior associated to $p_{a,b}$ is the space spanned by $(1 \mid 0, 0, \cdots)$. Note in particular that $\mathcal{B}(p_{a,b})$ is one-dimensional for *all* values of $(a,b) \neq (0,0)$, since the singularity that occurs at $a = 0$ for nonhomogeneous behaviors parametrized by degree-1 nonhomogeneous polynomials of the form $p(s) = as + 1$ is exactly "filled up". This is a special case of the main result of this paper, which states that there is a one-to-one connection between (i) homogeneous behaviors of degree n with m inputs and p outputs, (ii) triples of matrices (F,G,H) of sizes $n \times (n+m)$ (F,G) and $(p+m) \times (n+m)$ (H) satisfying certain minimality conditions, modulo similarity, and (iii) homogeneous polynomial matrices of size $p \times (p+m)$ with row degrees summing up to n, modulo left multiplication by homogeneous unimodular matrices.

The dehomogenization $P(s,1)$ of a homogeneous polynomial matrix $P(s,t)$ will be written simply as $P(s)$. By identifying polynomials with elements of $\mathcal{L}_+$, we can look at the matrix of polynomials $P(s)$ as a matrix with entries in $\mathcal{L}$, and so (using the multiplicative structure of $\mathcal{L}$) as a mapping from $\mathcal{L}^q$ to $\mathcal{L}^p$ where p is the number of rows of $P(s,t)$. Immediately from the definitions we now have the following characterization of $\mathcal{B}(P)$.

LEMMA 2.7 *Let the row degrees of the homogeneous polynomial $P(s,t)$ be $\nu_1, \cdots, \nu_p$. An element $w(s) \in \mathcal{L}^q$ belongs to $\mathcal{B}(P)$ if and only if the entries of the p-vector $P(s)w(s)$ are polynomials of degrees at most $\nu_1 - 1, \cdots, \nu_p - 1$ respectively.*

In the sequel we will associate to a homogeneous polynomial matrix $P(s,t)$ of size $p \times q$ and of row degrees $\nu_1, \cdots, \nu_p$ in a natural way a vector space of dimension $n := \sum_{i=1}^p \nu_i$. For this note that the set of p-vectors having the property that the i-th component is a homogeneous polynomial of degree $\nu_i - 1, i = 1, \ldots, p$ has in a natural way the structure of an $\mathbb{R}$-vector space. Obviously the dimension of this space is $\sum_{i=1}^p \nu_i = n$, the McMillan degree of the associated homogeneous polynomial $P(s,t)$. Since this vector space is closely related with the state space we will abbreviate it with X_ν. The analogous space of p-vectors whose i-th component is a homogeneous polynomial of degree ν_i will be denoted by $X_{\nu+1}$. The dimension of this space is $n+p$.

DEFINITION 2.8 A $p \times n$ matrix $X(s,t)$ whose columns form a $\mathbb{R}$-basis of the vector space X_ν will be called a *basis matrix* of size ν.

We define the *canonical basis matrix* $\hat{X}(s,t)$ as the matrix of size $p \times n$ given by

$$\hat{X}(s,t) = \begin{bmatrix} s^{\nu_1-1} & s^{\nu_1-2}t & \cdots & t^{\nu_1-1} & 0 & 0 & \cdots & \cdots & 0 \\ 0 & 0 & & 0 & s^{\nu_2-1} & \cdots & & & \vdots \\ \vdots & \vdots & & & & & \ddots & & 0 \\ 0 & 0 & \cdots & & & & & \cdots & t^{\nu_p-1} \end{bmatrix}. \tag{2.11}$$

REMARK 2.9 Every basis matrix has a unique description of the form $\hat{X}(s,t)S^{-1}$, where $S \in Gl_n$ is an $n \times n$ invertible matrix. In particular any two basis matrices are related to each other through a simple Gl_n transformation.

The result stated in the lemma above can now be reformulated as follows. Let $X(s,t)$ be any basis matrix and let $X(s) = X(s,1)$. Then one has that

$$\mathcal{B}(P) = \{w(s) \in \mathcal{L}^q \mid P(s)w(s) \in \operatorname{span}_{\mathbb{R}} X(s)\}.$$

It will be convenient to use polynomial representations that are of the above form but are not necessarily derived explicitly from a homogeneous polynomial matrix. For any pair of polynomial matrices $(R(s), V(s))$, where $R(s)$ and $V(s)$ have the same number of rows, we can define

$$\mathcal{B}(R,V) \;=\; \{w(s) \in \mathcal{L}^q \mid R(s)w(s) \in \operatorname{span}_{\mathbb{R}} V(s)\}. \tag{2.12}$$

Finally, a third representation that we shall use is the first-order representation. Consider a triple of real matrices (F, G, H) where F and G both have size $n \times (n+m)$ and H has size $q \times (n+m)$. With this triple we associate a behavior as follows:

$$\mathcal{B}(F,G,H) \;=\; \{w \in \mathcal{L}^q \mid w = Hz \text{ for some } z \in \mathcal{B}(sG - tF)\}. \tag{2.13}$$

Note that the operator associated to the homogeneous polynomial matrix $sG - tF$ is $\sigma_0(G - \tau F) = \sigma\tau_0(\sigma G - F)$. We may therefore also write the above definition in the form

$$\mathcal{B}(F,G,H) = \{w(s) \in \mathcal{L}^q \mid \exists z \in \mathcal{L}^{n+m},\; x_0 \in \mathbb{R}^n \text{ s.t. } (sG - F)z(s) = x_0 \text{ and } w(s) = Hz(s)\}.$$

We now turn to the relation between homogeneous behaviors and behaviors defined on $\mathbb{Z}_+$ such as for instance in [14]. The latter are given in 'AR representation' as follows. Let $P(s)$ be a polynomial matrix and let S denote the standard left shift that takes $(w_1, w_2, \cdots)$ to $(w_2, w_3, \cdots)$; then define $\mathcal{B}_-(P) = \{w \mid P(S)w = 0\}$. Identifying the space of one-sided sequences with $\mathcal{L}^q_-$, we get the following simple embedding of the set of 'standard' behaviors in the space of homogeneous behaviors.

PROPOSITION 2.10 *Let $P(s,t)$ be a homogeneous polynomial matrix, and let $P(s)$ be its dehomogenization. We have:*

$$\mathcal{B}_-(P) \;=\; \mathcal{B}(P) \cap \mathcal{L}^q_-. \tag{2.14}$$

PROOF Let w be a sequence in $\mathcal{B}_-(P)$. The fact that $P(S)w = 0$ means that $P(s)w(s)$ is polynomial. Moreover, it is immediate from the multiplication rule that the i-th entry of $P(s)w(s)$ is a polynomial of degree at most $\mu_i - 1$, where μ_i denotes the maximum of the degrees of the entries in the i-th row of $P(s)$. Since $\mu_i \le \nu_i$ where ν_i is the degree of the i-th row of $P(s,t)$, it follows that $w \in \mathcal{B}(P)$. Conversely, let w be a one-sided (strictly proper) sequence in $\mathcal{B}(P)$. Then we know that $P(s)w(s)$ is polynomial and so $P(S)w = 0$. □

To get a similar result for (R,V)-representations, we need to impose a condition involving the space X_R that is defined as follows [2, 8, 4]:

$$X_R \;=\; \{g(s) \in \mathcal{L}^p_+ \mid g(s) = R(s)w(s) \text{ for some } w(s) \in \mathcal{L}^q_-\}. \tag{2.15}$$

Using the same reasoning as above, we then obtain the following.

PROPOSITION 2.11 *Let $(R(s), V(s))$ be a pair of polynomial matrices having the same number of rows. If $X_R \subset \operatorname{span}_{\mathbb{R}} V(s)$, then*

$$\mathcal{B}_-(R) \;=\; \mathcal{B}(R,V) \cap \mathcal{L}^q_-. \tag{2.16}$$

A subspace of $\mathcal{L}^q$ may be given in image representation as $\operatorname{im} M(s)$ or in kernel representation as $\ker N(s)$, where $M(s)$ and $N(s)$ are matrices over $\mathcal{L}$. If $M(s)$ and $N(s)$ are rational matrices, we may also look at $\operatorname{im} M(s)$ and $\ker N(s)$ as curves, that is, as $\operatorname{Grass}(m,n)$- valued mappings defined almost everywhere (i. e. everywhere except for a finite number of points) on $\mathbb{C}$. The following lemma, which will be needed below, essentially says that these two points of view are equivalent.

LEMMA 2.12 *Let $M(s)$ and $N(s)$ be rational matrices. In this case we have $\operatorname{im}_{\mathbb{C}} M(s) = \ker_{\mathbb{C}} N(s)$ for almost all $s \in \mathbb{C}$ if and only if $\operatorname{im}_{\mathcal{L}} M(s) = \ker_{\mathcal{L}} N(s)$.*

PROOF We may assume without loss of generality that $M(s)$ and $N(s)$ have full column rank and full row rank respectively as matrices over $\mathbb{R}(s)$ (or, equivalently, as matrices over $\mathcal{L}$). Let $M(s)$ have size $q \times m$ and let $N(s)$ have size $p \times q$. If $\operatorname{im} M(s) = \ker N(s)$ for almost all $s \in \mathbb{C}$, then $N(s)M(s) = 0$

almost everywhere and hence everywhere on $\mathbb{C}$. This shows that $\mathrm{im}_{\mathbb{C}}\, M(s) \subset \mathrm{im}_{\mathbb{C}}\, N(s)$. From $\mathrm{im}\, M(s) = \ker N(s)$ it also follows that $p+m=q$, and together with the full rank assumption this implies that $\mathrm{im}_{\mathbb{C}}\, M(s) = \ker_{\mathbb{C}} N(s)$. The converse is obtained by a similar argument. □

The following lemma is standard; we provide a proof for completeness.

LEMMA 2.13 *Let A, B, X, and Y be matrices such that $AX+BY=0$. If $[A \;\; B]$ and X have full row rank, then also B has full row rank.*

PROOF Let η be a row vector such that $\eta B = 0$. It follows from $\eta(AX+BY)=0$ that $\eta AX = 0$ and consequently $\eta A = 0$ because X has full row rank. But then we have $\eta[A \;\; B] = 0$, which implies that $\eta = 0$. □

The result in the lemma below is standard as well; it is related for instance to the fact that the observability indices and the controllability indices of a linear system both sum up to the same value, which coincides with the state space dimension n. We shall need the following form. By the *degree* of a polynomial matrix of full row or column rank, we mean the maximum of the degrees of its full-size minors.

LEMMA 2.14 *If $P(s)$ and $Q(s)$ are polynomial matrices of full column rank and full row rank respectively and we have*

$$\ker Q(s) \;=\; \mathrm{im}\, P(s) \tag{2.17}$$

for all $s \in \mathbb{C}$, then the degree of $P(s)$ is equal to the degree of $Q(s)$.

PROOF From the fact that the equality 2.17 holds for all s it follows that the Smith forms of $Q(s)$ and $P(s)$ must be $[I \mid 0]$ and $[{I \atop 0}]$ respectively. Therefore we can find (see for instance [7, p. 382]) a polynomial matrix $T(s)$ such that $[{T(s) \atop Q(s)}]$ is unimodular. Define $Z(s)$ by

$$\begin{bmatrix} T(s) \\ Q(s) \end{bmatrix} P(s) \;=\; \begin{bmatrix} Z(s) \\ 0 \end{bmatrix};$$

it follows that $Z(s)$ is unimodular. Without loss of generality, we may assume that $Q(s) = [Q_1(s) \mid Q_2(s)]$ where $Q_1(s)$ gives a minor with the maximal degree. Partitioning $T(s)$ and $P(s)$ accordingly, we get

$$\begin{bmatrix} T_1(s) & T_2(s) \\ Q_1(s) & Q_2(s) \end{bmatrix} \begin{bmatrix} P_1(s) & 0 \\ P_2(s) & I \end{bmatrix} \;=\; \begin{bmatrix} Z(s) & 0 \\ 0 & Q_2(s) \end{bmatrix}$$

which shows that $\det P_1(s) = c \cdot \det Q_1(s)$ for some nonzero constant c. It follows that the degree of $P(s)$ must be at least as large as the degree of $Q(s)$. Since a similar reasoning provides the reverse inequality, the lemma is proved. □

3 Realization of homogeneous systems

Since we have associated homogeneous behaviors both to triples (F,G,H) and to homogeneous polynomial matrices $P(s,t)$, we can now state the following definition.

DEFINITION 3.1 A triple of constant matrices (F,G,H) is said to be a *realization* of the polynomial matrix $P(s,t)$ if $\mathcal{B}(F,G,H) = \mathcal{B}(P)$.

The following lemma gives a sufficient condition for a triple (F,G,H) to be a realization of a homogeneous polynomial matrix $P(s,t)$.

LEMMA 3.2 *Let $P(s,t)$ be a $p \times q$ homogeneous polynomial matrix with row degrees $\nu = (\nu_1, \ldots, \nu_p)$. Let $X(s,t)$ be a spanning matrix for the vector space X_ν. If a triple (F,G,H) is such that the equality*

$$\mathrm{im}\begin{bmatrix} sG-tF \\ H \end{bmatrix} = \ker\,[-X(s,t) \mid P(s,t)] \tag{3.1}$$

holds for almost all $(s,t)\in\mathbb{C}^2\setminus\{(0,0)\}$, *then* $\mathcal{B}(F,G,H)$ *is equal to* $\mathcal{B}(P)$.

PROOF First take $w\in\mathcal{B}(F,G,H)$, and let $z(s)$ and the constant vector x_0 be such that $w(s)=Hz(s)$ and $(sG-F)z(s)=x_0$. We then have, by (3.1) and lemma 2.12,

$$\begin{bmatrix} x_0 \\ w(s) \end{bmatrix} \in \mathrm{im}_{\mathcal{L}}\begin{bmatrix} sG-F \\ H \end{bmatrix} = \ker_{\mathcal{L}}\,[-X(s)\mid P(s)].$$

It follows that $P(s)w = X(s)x_0 \in \mathrm{span}_{\mathbb{R}}\,X(s)$ and so $w\in\mathcal{B}(P)$. For the reverse inclusion, take $w\in\mathcal{B}(P)$. Then we must have $P(s)w(s)=X(s)x_0$ for some constant x_0 and by the formula above we get $w\in\mathcal{B}(F,G,H)$. □

The following theorem states that every polynomial matrix has a realization. Moreover, it is shown that the matrices (F,G,H) in the realization can be chosen to satisfy certain requirements that will later be seen to be minimality conditions. Our proof is based on an elementary realization algorithm which applies *mutatis mutandis* (compare with [13]) to standard (non-homogeneous) systems as well. This algorithm was first announced in [12]. Though we formulate the proof for real homogeneous systems $P(s,t)$ we would like to remark that the same proof is also valid for homogeneous matrices $P(s,t)$ which are defined over an arbitrary base field.

THEOREM 3.3 *Every homogeneous system* $P(s,t)$ *has a realization* (F,G,H) *satisfying the following properties:*

1. $\mathrm{rank}\begin{bmatrix} sG-tF \\ H \end{bmatrix} = m+n$ *for all* $(s,t)\in\mathbb{C}^2\setminus\{(0,0)\}$.
2. $\mathrm{rank}_{\mathbb{R}(s)}\,[sG-tF]=n$.

Finally $P(s_0,t_0)$ *has rank* p *if and only if* s_0G-t_0F *has rank* n, *in particular* $P(s,t)$ *is controllable if and only if* $sG-tF$ *is controllable.*

PROOF The existence of a realization is shown via a simple realization algorithm. Let $\nu=(\nu_1,\ldots,\nu_p)$ be the set of row degrees of $P(s,t)$ and let $X(s,t)$ be a basis matrix for the vector space X_ν.

Algorithm: Consider the $p\times(2n+m+p)$ matrix

$$A(s,t) := [tX(s,t)\mid -sX(s,t)\mid P(s,t)]\,.$$

By definition the i-th row of $A(s,t)$ contains homogeneous polynomials of degree ν_i and so $A(s,t)$ defines a linear map from $\mathbb{R}^{2n+m+p}$ to the vector space $X_{\nu+1}$. Note that $\dim X_{\nu+1} = n+p$, and that the mapping defined by $A(s,t)$ is surjective (since $X(s,t)$ is a basis matrix for X_ν, and every polynomial $x(s,t)$ in $X_{\nu+1}$ can be written as $x(s,t)=tx_1(s,t)+sx_2(s,t)$ for some $x_1(s,t)$ and $x_2(s,t)$ in X_ν). Therefore the kernel of this mapping is an $(n+m)$-dimensional subspace of $\mathbb{R}^{2n+m+p}$. Let an image representation for this subspace be given by the matrix

$$\begin{bmatrix} F \\ G \\ H \end{bmatrix} \tag{3.2}$$

where the partitioning corresponds to the partitioning of $A(s,t)$. It is immediate from Lemma 3.2 that the triple (F,G,H) defines a realization of $P(s,t)$.

We now show that the triple (F,G,H) constructed this way does have the properties 1. and 2. as claimed in the theorem. To show 1., we have to prove that

$$\begin{bmatrix} s_0G - t_0F \\ H \end{bmatrix} z_0 \;=\; 0 \tag{3.3}$$

for some $(s_0, t_0) \in \mathbb{C}^2 \setminus \{(0,0)\}$ and $z_0 \in \mathbb{R}^{n+m}$ implies that $z_0 = 0$. So suppose that (3.3) holds. Then we have in particular that $s_0Gz_0 = t_0Fz_0$, so there must exist an $x_0 \in \mathbb{R}^n$ and constants α and β such that $Gz_0 = \alpha x_0$ and $Fz_0 = \beta x_0$. We then have $X(s,t)(sG - tF)z_0 = X(s,t)(\alpha s - \beta t)x_0$. On the other hand we also have $X(s,t)(sG - tF)z_0 = P(s,t)Hz_0 = 0$, and so it follows that $x_0 = 0$ so that $Fz_0 = Gz_0 = 0$. Since also $Hz_0 = 0$, it follows from the independence of the columns of the matrix in (3.2) that $z_0 = 0$.

It remains to be shown that $\operatorname{rank}_{\mathbb{R}(s)}[sG - tF] = n$. This will be a trivial consequence of the controllability part which we will now prove.

Controllability: Take $(s_0, t_0) \neq (0,0)$. It follows from the identity $P(s,t)H = X(s,t)(sG - tF)$ that $Hz \in \ker P(s_0,t_0)$ for all $z \in \ker(s_0G - t_0F)$. Moreover, it follows from property 1. that $\ker H \cap \ker(s_0G - t_0F) = \{0\}$, so that

$$\dim\ker P(s_0,t_0) \geq \dim\ker(s_0G - t_0F) \geq m \tag{3.4}$$

where the final equality is obvious from the fact that $sG - tF$ has size $n \times (n+m)$. Since $P(s,t)$ has size $p \times (p+m)$, it follows that $s_0G - t_0F$ must have full row rank when $P(s_0,t_0)$ has full row rank. The converse is immediate from Lemma 2.13. □

REMARK 3.4 For a fixed basis matrix $X(s,t)$ the coefficients of the realization (F,G,H) can be chosen continuously dependent on the coefficients of the homogeneous system $P(s,t)$. This follows from the presented realization algorithm, in particular identity (3.2). A similar result using the so called 'Fuhrmann realization' [2, Ch. I.10] has been established by Glüsing-Lüerßen [5].

In addition to this realization theorem, which describes the transformation from polynomial to state space form, we shall also need a result that produces a polynomial representation starting from a state space description. Such a result may be called an 'elimination theorem', since essentially what is involved is the elimination of the state variables.

THEOREM 3.5 *For any triple (F,G,H) satisfying conditions 1. and 2. of Thm. 3.3, there exists a pair of homogeneous polynomial matrices $(X(s,t), P(s,t))$ such that the following holds:*

(i) $\ker[-X(s,t) \mid P(s,t)] = \operatorname{im}\begin{bmatrix} sG-tF \\ H \end{bmatrix}$ *for all $(s,t) \neq (0,0)$*

(ii) $P(s,t)$ has full row rank

(iii) the columns of $X(s,t)$ form a basis for the vector space X_ν, where $\nu = (\nu_1, \ldots, \nu_p)$ is the set of row degrees of $P(s,t)$.

In particular, we have $\mathcal{B}(F,G,H) = \mathcal{B}(P)$.

PROOF We dehomogenize with respect to a point (s_0,t_0) such that $s_0G - t_0F$ has full row rank. For ease of notation and without loss of generality we shall assume that we can take $(s_0,t_0) = (1,0)$, so that the matrix G has full row rank. By any one of a variety of methods (see for instance [7, p. 488] or [1, p. 61]), one can find polynomial matrices $X(s)$ and $P(s)$ such that

$$\ker[-X(s) \mid P(s)] \;=\; \operatorname{im}\begin{bmatrix} sG - F \\ H \end{bmatrix} \tag{3.5}$$

for all complex s. Since we can premultiply $X(s)$ and $P(s)$ by a unimodular matrix without affecting the above property, we may assume that the matrix $[-X(s) \mid P(s)]$ is row reduced. Let n be the number of rows of G. Because $sG - tF$ has full row rank n, it follows from the relation (3.5) and

Lemma 2.14 that the sum of the row degrees of $[-X(s) \mid P(s)]$ must be equal to n. Let the row degrees of $P(s)$ be denoted by $\nu_1, \ldots, \nu_p$. Because G has full row rank and $X(s)(sG-F) = P(s)H$, the row degrees of $X(s)$ must be strictly less than those of $P(s)$. It follows that the ν_i are also the row degrees of $[-X(s) \mid P(s)]$ so that they must sum up to n. It follows that the columns of $X(s)$ all belong to the vector space X_ν. Since $X(s)$ has n columns and the columns are independent by Lemma 3.1 in [3], it follows that actually $X(s)$ must be a basis matrix. Finally note that $P(s)$ must have full row rank by Lemma 2.13.

Now let $X(s,t)$ and $P(s,t)$ be the homogenizations of $X(s)$ and $P(s)$ respectively. Property (i) in the statement of the theorem follows from (3.5) together with our assumptions that $[-X(s) \mid P(s)]$ is row reduced and that $[\begin{smallmatrix} G \\ H \end{smallmatrix}]$ has full column rank. Properties (ii) and (iii) are immediate from what has been said above. The final claim is immediate from Lemma 3.2. $\square$

4 Homogeneous systems and their homogeneous behaviors

We shall need the following uniqueness theorem for polynomial representations (compare [3, Thm. 3.10]).

Theorem 4.1 *Let $(R_1(s), V_1(s))$ and $(R_2(s), V_2(s))$ be two pairs of polynomial matrices, and assume that for $i = 1, 2$ the following holds:*

(i) $R_i(s)$ has full row rank,

(ii) $\operatorname{span}_{\mathbb{R}} V_i(s) \supset X_{R_i}$,

(iii) $V_i(s)$ has full column rank over $\mathbb{R}$.

Under these conditions, we have $\mathcal{B}(R_1, V_1) = \mathcal{B}(R_2, V_2)$ if and only if there exists a unimodular polynomial matrix $U(s)$ and a nonsingular constant matrix S such that $R_2(s) = U(s)R_1(s)$ and $V_2(s) = U(s)V_1(s)S$.

Proof The 'if' part is immediate from the definition. For the converse, first note that we must have $\mathcal{B}_-(R_1) = \mathcal{B}_-(R_2)$ by Prop. 2.10. It then follows from the uniqueness theorem for polynomial representations of behaviors on $\mathbb{Z}_+$ that there must exist a unimodular matrix $U(s)$ such that $R_2(s) = U(s)R_1(s)$. So for the rest of the proof we may assume that this unimodular transformation has already been carried out, and we write $R_1(s) = R_2(s) = R(s)$. Since both $V_1(s)$ and $V_2(s)$ have full column rank over $\mathbb{R}$, it will be sufficient to show that $\operatorname{span}_{\mathbb{R}} V_1(s) = \operatorname{span}_{\mathbb{R}} V_2(s)$; by symmetry, is is actually already sufficient to prove that $\operatorname{span}_{\mathbb{R}} V_1(s) \subset \operatorname{span}_{\mathbb{R}} V_2(s)$.

Take $g(s) \in \operatorname{span}_{\mathbb{R}} V_1(s)$. Because $R(s)$ has full row rank, there is a $w(s)$ such that $g(s) = R(s)w(s)$. Write $w(s) = w_+(s) + w_-(s)$ where $w_+(s) \in \mathcal{L}_+^q$ and $w_-(s) \in \mathcal{L}_-^q$. Because $R(s)w_-(s) = g(s) - R(s)w_+(s)$ is a polynomial, we have that $R(s)w_-(s) \in X_R$ and so, by assumption (ii) above, $R(s)w_-(s)$ belongs to both $\operatorname{span}_{\mathbb{R}} V_1(s)$ and $\operatorname{span}_{\mathbb{R}} V_2(s)$. Since also $R(s)w(s) \in \operatorname{span}_{\mathbb{R}} V_1(s)$, we get that $R(s)w_+(s) \in \operatorname{span}_{\mathbb{R}} V_1(s)$ and so $w_+(s) \in \mathcal{B}(R, V_1) = \mathcal{B}(R, V_2)$. Now we have both $R(s)w_+(s) \in \operatorname{span}_{\mathbb{R}} V_2(s)$ and $R(s)w_-(s) \in \operatorname{span}_{\mathbb{R}} V_2(s)$, so $g(s) = R(s)w(s) \in \operatorname{span}_{\mathbb{R}} V_2(s)$. $\square$

The crucial property that we have been aiming for is stated in the theorem below.

Theorem 4.2 *If two triples (F_1, G_1, H_1) and (F_2, G_2, H_2) both satisfy properties 1. and 2. of Thm. 3.3, then $\mathcal{B}(F_1, G_1, H_1) = \mathcal{B}(F_2, G_2, H_2)$ if and only if there exist constant nonsingular matrices S and T such that $F_2 = SF_1T^{-1}$, $G_2 = SG_1T^{-1}$, and $H_2 = H_1T^{-1}$.*

Proof By the elimination theorem of the previous section, we can find for $i = 1, 2$ homogeneous polynomial matrices $X_i(s,t)$ and $P_i(s,t)$ satisfying conditions (i)–(iii) of that theorem. In particular it follows that $\mathcal{B}(P_1, X_1) = \mathcal{B}(F_1, G_1, H_1) = \mathcal{B}(F_2, G_2, H_2) = \mathcal{B}(P_2, X_2)$. The assumptions of the uniqueness theorem Thm. 4.1 are satisfied and so we can conclude that there exists a unimodular matrix $U(s)$ and a constant nonsingular matrix S such that $P_2(s) = U(s)P_1(s)$ and $X_2(s) = U(s)X_1(s)S$. It follows that

$$\operatorname{im}\begin{bmatrix} sG_2 - F_2 \\ H_2 \end{bmatrix} = \operatorname{im}\begin{bmatrix} sSG_1 - SF_1 \\ H_1 \end{bmatrix} \tag{4.1}$$

and by our assumptions on both triples this implies the results claimed in the theorem (see the proof of Thm. 4.1 in [3] for details). □

REMARK 4.3 It follows as in [3, Thm. 4.2] that the conditions 1. and 2. of Thm.3.3 are actually minimality conditions. To be precise, we have the following: if $\mathcal{B}(F_1, G_1, H_1) = \mathcal{B}(F_2, G_2, H_2)$ where $sG_1 - tF_1$ has size $n_1 \times (n_1 + m_1)$ and $sG_2 - tF_2$ has size $n_2 \times (n_2 + m_2)$, and the triple (F_1, G_1, H_1) satisfies conditions 1. and 2., then $n_2 \geq n_1$ and $m_2 \geq m_1$. So it is justified to refer to triples (F, G, H) satisfying conditions 1. and 2. as *minimal* triples.

REMARK 4.4 The main result in [11] states that there is a one-to-one relation between minimal triples modulo similarity equivalence and full row rank homogeneous polynomial matrices modulo left multiplication by homogeneous unimodular matrices. So from the above theorem we also obtain a one-to-one connection between the latter quotient space and homogeneous behaviors.

5 CONCLUSIONS

Using the set of formal Laurent series as a universum we introduced a new class of behaviors which we called homogeneous behaviors. Every homogeneous behavior can be described either through a homogeneous polynomial matrix or through a triple of matrices (F, G, H) inducing a generalized first-order representation of the homogeneous behavior. The relation between higher-order and first-order representation was explained. Both the equivalence classes of first-order representations and the equivalence classes of higher-order representations can be seen as points in a certain quotient scheme due to Grothendieck (cf. [11]), and therefore from a mathematical point of view our results come down to giving these points a concrete interpretation as elements in the Grassmannian of the space of vector-valued formal Laurent series.

REFERENCES

[1] F.M. CALLIER AND C.A. DESOER (1982). *Multivariable Feedback Systems.* Springer, New York.

[2] P. A. FUHRMANN (1981). *Linear Systems and Operators in Hilbert Space.* McGraw-Hill, New York, NY.

[3] A. H. W. GEERTS AND J. M. SCHUMACHER (1994). Impulsive-smooth behavior in multimode systems. Part II: Minimality and equivalence. Report BS-R9432, CWI, Amsterdam, to appear in *Automatica.*

[4] A. H. W. GEERTS AND J. M. SCHUMACHER (1996). Impulsive-smooth behavior in multimode systems. Part I: State-space and polynomial representations. *Automatica* 32, 747–758.

[5] H. GLÜSING-LÜERSSEN (1995). Continuous state representations for AR systems. *Math. Contr., Sign., and Syst.* 8, No. 1, 82–95.

[6] M. L. J. HAUTUS AND M. HEYMANN (1978). Linear feedback—an algebraic approach. *SIAM J. Control* 16, 83–105.

[7] T. KAILATH (1980). *Linear Systems.* Prentice-Hall, Englewood Cliffs, N.J.

[8] M. KUIJPER AND J. M. SCHUMACHER (1990). Realization of autoregressive equations in pencil and descriptor form. *SIAM J. Control Optim.* 28, No. 5, 1162–1189.

[9] V. G. LOMADZE (1990). Finite-dimensional time-invariant linear dynamical systems: algebraic theory. *Acta Appl. Math* 19, 149–201.

[10] M. S. RAVI AND J. ROSENTHAL (1994). A smooth compactification of the space of transfer functions with fixed McMillan degree. *Acta Appl. Math.* 34, 329–352.

[11] M. S. RAVI AND J. ROSENTHAL (1995). A general realization theory for higher order linear differential equations. *Systems & Control Letters* 25, No. 5, 351–360.

[12] M. S. RAVI, J. ROSENTHAL AND J. M. SCHUMACHER (1995). A realization theory for homogeneous AR-systems, an algorithmic approach, in *Proc. IFAC Conference on System Structure and Control* (Nantes, France, July 1995), 183–188.

[13] J. ROSENTHAL AND J. M. SCHUMACHER. Realization by inspection. CWI Report BS-R9534, 1995.

[14] J. C. WILLEMS (1986). From time series to linear system. Part I: Finite dimensional linear time invariant systems. *Automatica* 22, 561–580.

[15] J. C. WILLEMS (1989). Models for dynamics. *Dynamics Reported* 2, 171–269.

[16] J. C. WILLEMS (1991). Paradigms and puzzles in the theory of dynamical systems. *IEEE Trans. Automat. Control* AC-36, No. 3, 259–294.